

# FACT SHEET

---

## Tobacco Use and Policy

---

### Estill County

|   | Estill County | Kentucky |
|---|---------------|----------|
| Adult Smoking <sup>1</sup> , 2005-2007  | 17.1          | 27.4     |
| Adult Quit Attempts <sup>1</sup> , 2005-2007  | 33.3          | 48.8     |
| Adult Participation in Cessation <sup>2</sup> , 2008  | *             | 83.9     |
| Smoking During Pregnancy <sup>3</sup> , 2006  | 38.1          | 26.5     |
| Manufacturers that Permit Indoor Smoking <sup>4</sup> , 2008  | 0.0           | 30.0     |
| Manufacturers that Offer Cessation <sup>4</sup> , 2008  | 100.0         | 66.0     |
| Manufacturers that Reimburse for Cessation <sup>4</sup> , 2008  | 100.0         | 59.4     |
| Percentage of Manufacturing Workers Covered by Indoor Smoking Policy <sup>4</sup> , 2008                          | 100.0         | 73.2     |
| Schools that Ban Smoking on School Grounds for Employees <sup>4</sup> , 2007                                      | 0.0           | 52.1     |
| Schools that Offer Cessation to Students and/or Employees <sup>4</sup> , 2007                                     | 0.0           | 33.5     |
| Schools that Post No Tobacco Signs <sup>4</sup> , 2007  | 50.0          | 73.2     |
| Percentage of Residents Covered by Comprehensive Smoke-free Workplace Ordinance or Regulation <sup>5</sup> , 2009 | 0.0           | 29.9     |
| Kentucky Institute of Medicine Ranking <sup>6</sup> , 2007<br>(Higher numbers indicate poorer health outcomes)    | 89.0          | 60.5     |

\*Data unavailable.

<sup>1</sup>Unweighted, 3-year pooled estimate--- Source: Kentucky Behavioral Risk Factor Surveillance System, 2005-2007;

Use caution interpreting this estimate for the county since the sample size is < 50.

<sup>2</sup>Rate per 10,000 adult smokers--- Source: Local Health Department Tobacco Cessation Survey, 2007. The state rate excludes data from Breathitt County since the number of smokers reached by tobacco cessation programs was significantly higher than all other health department service areas and including it would have made the state average artificially high.

<sup>3</sup>Kentucky Department for Public Health, Office of Vital Statistics, 2006. Data are considered preliminary. Use caution when interpreting this estimate.

<sup>4</sup>UK Tobacco Policy Research Program, <http://www.mc.uky.edu/tobaccopolicy/KentuckyDataReports/> --- Workplace policy 2008; School Policy 2007. Workplace policy study includes only manufacturers with 50 or more employees. School policy study includes public and private middle and high schools.

<sup>5</sup>UK Tobacco Policy Research Program, Percent Covered Database, January 2009: accessed from <http://www.mc.uky.edu/tobaccopolicy/Ordinances/Reports>.

<sup>6</sup>Kentucky Institute of Medicine. The Health of Kentucky: A County Assessment. Lexington, KY. 2007. Higher numbers indicate poorer health outcomes. A full report can be found at: <http://kviom.org/healthy2007a.pdf>.

**Prepared by:**

**Kentucky Center for Smoke-Free Policy**

**315 CON Bldg., 751 Rose St.**

**Lexington, KY 40536-0232**

**859-323-4587**

[www.kcsp.uky.edu](http://www.kcsp.uky.edu)

February 2009